

A Systematic Study on the Marine Sponges in Korea. 7. Demospongiae and Hexactinellida

Sim, Chung Ja and Kim, Mi Hee

(Department of Biology, Han Nam University, Daejeon 300-791, Republic of Korea)

한국산 해산해면류의 계통분류학적 연구 7. 보통해면류와 육방해면류

심 정 자 · 김 미 희
(한남대학교 생물학과)

적 요

1985년부터 1987년에 걸쳐 남한의 동·서·남해로부터 채집된 해면동물과 그동안 미해결로 보류되어 있던 표본들을 조사한 결과 총 24종이 밝혀졌는데 이 중에서 23종은 보통해면류 이었으며 6종이 한국미기록 종이고, 1종은 육방해면 동물로서 신종임이 판명되어 *Aphrocallistes jejuensis*라고 명명기재한다. 한국 미기록종에 대하여는 특기와 도판을 첨가하였다.

Key words: Systematic, Demospongiae, Hexactinellida, Korea.

INTRODUCTION

On the systematic study of Korean Demospongiae and Hexactinellida, 141 species were recorded from Korean waters (Kim *et al.*, 1968; Rho & Sim, 1972b; Sim, 1982). Of which, 139 species were Demospongiae and only two species were Hexactinellida (Rho & Sim, 1972a; Sim, 1982).

The present study is based on the materials collected from 17 localities (see Fig. 1) during the period from 1985 to 1987, and some preserved specimens were used for this experiment. The species identified consist of 24 species, 20 genera and 17 families. Of these species, one Hexactinellida from Pömsöm, Cheju-do is new to science. Six Demospongiae are unrecorded ones from Korea.

본 연구는 한국학술진흥재단의 “’86 첨단과학 기초연구지원”에 의한것임.




Fig. 1. A map showing the localities where the materials were collected. 1, Ulreungdo; 2, Mip'o; 3, Haeundae; 4, Pijindo; 5, Ch'ungmu; 6, Mijo-ri; 7, Sangju-ri; 8, Namhaedo; 9, Tölsando; 10, U-od; 11, Söngsanp'o; 12, Sögwip'o; 13, Pömsöm; 14, 33°00'00"N, 217°30'00"E; 15, Ch'ujado; 16, Manlip'o; 17, Anhung.

LIST OF SPECIES

Class Demospongiae

Subclass Ceractionomorpha

Order Halichondria

Family Hymeniacionidae De Laubenfels, 1936

1. *Hymeniacion sinapium* De Laubenfels, 1930

Family Halichondriidae Gray, 1867

2. *Halichondria okadai* (Kadota, 1922)

Family Ophlitaspongiidae De Laubenfels, 1936

3. * *Ophlitaspongia minor* Burton, 1959
4. *Ophlitaspongia noto* Tanita, 1963
5. * *Oxymycale paradoxa* De Laubenfels, 1935
6. * *Clathria madrepora* Dendy, 1921
7. *Clathria spinispicula* Tanita, 1968

Family Microciona Henschel, 1923

8. *Microciona longistyla* Burton, 1959

Family Myxillidae Henschel, 1923

9. * *Myxilla productus* Hoshino, 1981
10. *Myxilla setoensis* Tanita, 1961
11. *Lissodendoryx isodictyalis* (Carter, 1882)

Family Tedaniidae Ridley & Dendy, 1887

12. *Iotrochota baculifera* Ridley, 1884

Order Haplosclerida Topsent, 1928

Family Halicionidae De Laubenfels, 1932

13. *Haliclona permollis* (Bowerbank, 1866)

Family Callyspongiidae De Laubenfels, 1936

14. *Callyspongia elegans* (Thiele, 1899)
15. *Callyspongia elongate* (Ridley & Dendy, 1886)

Family Renieridae Schmidt, 1870

16. * *Gellius edaphus* De Laubenfels, 1930

Order Keratosa Grant, 1861

Family Spongiidae Gray, 1867

17. *Spongia officinalis* Linne, 1759

Subclass Tetractinomorpha

Order Epipolasida Sollas, 1888

Family Tethyidae Gray, 1867

18. *Tethya amamensis* Thiele, 1898

Order Hadromerida Tospent, 1894

Family Suberitidae Ridley et Dendy, 1886

19. *Suberites excellens* (Thiele, 1898)

Order Spirophorida Levi, 1956

Family Tetillidae Sollas, 1886

20. *Tetilla ovata* Thiele, 1898

Order Axinellida Bergquist, 1970

Family Raspailiidae Henschel, 1912

21. *Raspailia hirsuta* Thiele, 1898

Order Choristida Sollas, 1880

Family Kaliapsidae De Laubenfels, 1936

22. *Discodermia calyx* Döderlein, 1883

Family Stelletidae Carter, 1875

23. * *Stelletta validissima* Thiele, 1898

Class Hexactinellida

Order Hexasterophora F.E.Sch., 1899

Family Aphrocallistidae Gray, 1858

24. * *Aphrocallistes jejuensis*, nov. sp.

The asterisks(*) indicate the species which were newly recorded from Korean waters.

Family Hymeniacionidae De Laubenfels, 1936

Genus *Hymeniacion* Bowerbank, 1864

1. *Hymeniacion sinapium* De Laubenfels, 1930

Hymeniacion sinapium: De Laubenfels, 1932 (pp. 57-60, Text-fig. 29); Sim, 1985 (p. 6, pl. 1, figs. 5-6).

Halichondria japonika: Kim *et al.*, 1968 (p. 39, pl. 1, fig. 5, text-fig. 6); Rho *et al.*, 1969 (p. 155); Hoshino, 1970 (p. 22), 1971 (p. 23); Rho & Sim, 1972 a (pp. 183-184), 1972 b (p. 126); Hoshino, 1974, (p. 10), 1976 (p. 6); Sim, 1982 (p. 192).

Material examined: Söngsanp'o, VII/1985; Sögwip'o, VII/1987; Tölsando VIII/1987.

Distribution: California, Japan, Korea (East, South, West).

Family Halichondriidae Gray, 1867

Genus *Halichondria* Fleming, 1928

2. *Halichondria okadai* (Kadota, 1922)

Halichondria okadai: Tanita, 1964 (p. 18, pl. 1, fig. 6); Kim *et al.*, 1968 (p. 39, pl. 1, fig. 6, text-fig. 7); Hoshino, 1970 (p. 22), 1971 (p. 23), 1974 (p. 10).

Material examined: Mijo-ri, VII/1983; Söngsanp'o, VII/1985.

Distribution: Korea (Korea strait, Cheju Island), Japan.

Family Ophlitaspongidae De Laubenfels, 1936

Genus *Ophlitaspongia* Bowerbank, 1964

3. * *Ophlitaspongia minor* Burton, 1959

(pl. 1, figs. 1-2)

Ophlitaspongia minor Burton 1959 (pp. 246-247, text-fig. 27).

Material examined: Sögwip'o, VII/1982.

Remarks: Sponge is long, irregularly branched, 5-6mm in diameter, 17cm in length. The texture is firm and the surface of the body is a little velvety with the protruding spicules. Colour in alcohol beige.

Spicules:

Large Style	439 - 732 × 25-27 μm.
Small Style	333 - 386 × 20-23 μm.
Subtylostyle	306-704 × 4-8 μm.
Toxa	40-119 μm.
Isochela	16-20 μm.

Distribution: Korea, Arabian Sea.

4. *Ophlitaspongia noto* Tanita, 1963

Ophlitaspongia noto Tanita, 1963 (pp. 124-125, pl. 4, fig. 3, text-fig. 3), 1964 (pp. 17-18, pl. 1, fig. 4), 1965 (p. 48); Kim: *et al.*, 1968 (p. 41, pl. 3, fig. 12, text-fig. 13); Rho *et al.*, 1969 (p. 158); Rho & Sim, 1972a (p. 185), 1972b (p. 128); Hoshino, 1981 (p. 173); Sim, 1982 (P. 195).

Material examined: Sŏgwip'o, VII/1984.

Distribution: Korea, Japan.

Genus *Oxymycale* Hentschel, 1929

5. * *Oxymycale paradoxa* De Laubenfels, 1935 (pl. 4, figs. 1-2)

Oxmycale paradoxa De Laubenfels, 1935 (p. 5, fig. 2).

Material examined: 33° 00'00"N, 127°30'00"E, 145m depth, IV/1987.

Remarks: This Sponge is hemispherical, size 5 × 3cm. On the top occurs oscule, 0.6mm in diameter. Texture is compressible and soft. Colour in spirit is dirty grey.

Spicules:

Style	501-714 × 8-14μm.
Oxea	399-728 × 4-17μm.
Anisochela	84-115μm.
Sigma	21-70μm.
Raphid	70-84μm.

Distribution: Mexico, Korea.

Genus *Clathria* Schmidt, 1862

6. * *Clathria madrepora* Dendy, 1921 (p 1.2, figs. 1-2)

Clathria madrepora: Dendy, 1921 (pp. 68-69, pl. 5, fig. 3, pl. 14, figs. 1a-d).

Material examined: Sŏgwip'o VII/1982.

Remarks: Sponge ramified, thin noodle like irregularly branched, averaging about 3mm in diameter. Pore and oscules not conspicuous. Texture is tough. Colour in spirit pale brown.

Spicules:

Large tylostyle	254-308 × 7-14μm.
Small tylostyle	154-224 × 4-7μm.
Acantho subtylostyle	91-112 × 7-10μm.
Isochela	12-14μm.

Distribution: Korea, Seychelles.

7. *Clathria spinispicula* Tanita, 1968

Clathria spinispicula Tanita, 1968 (pp. 48-49. pl. 1, fig. 6, text-fig. 8).

Clathria spinispicula: Rho & Sim, 1972 a (p. 185, pl. 4, figs. 9-10); Hoshino, 1981 (p. 161); Sim, 1982 (p. 193, pl. 6, fig. 3).

Material examined: Sŏgwip'o, VII/1982.

Distribution: Korea, Japan.

Family Microcionidae Henschel, 1923

Genus *Microciona* Bowerbank, 1864

8. *Microciona longistyla* Burton, 1959

Microciona longistyla Burton, 1959 (pp. 249-250, text-fig. 29); Rho & Sim, 1976b (p. 100, pl. 2, figs. 7-8).

Material examined: Sŏgwip'o, VII/1982.

Distribution: Korea, South Arabian Sea (?).

Family Myxillidae Hentschel, 1923

Genus *Myxilla* Schmidt, 1862

9. * *Myxilla productus* Hoshino, 1981

(pl.3, figs. 1-2)

Myxilla productus Hoshino, 1981 (pp. 138-140, pl. 6, fig. 2, text-fig. 58).

Material examined: Sŏgwip'o, VII/1982.

Remarks: Sponge is long, Solid cylindrical branched, 25cm in height, 2cm wide, 8-10mm in thickness. Texture hard and tough. Colour in alcohol is ivory.

Spicules:

Large acanthostyle	252-294 × 11-17µm.
Small acanthostyle	100-157 × 6-14µm.
Tornote	168-196 × 7-12µm.
Sigma	56-70µm.
Isochela	28-39µm.
Biotulate	12-15µm.

Distribution: Japan, Korea.

10. *Myxilla setoensis* Tanita, 1961

Myxilla setoensis Tanita, 1961 (pp. 158-160, pl. 2, figs. 8-9, text-fig. 3); Kim *et al.*, 1968 (p. 40, pl. 2, fig. 10, text-fig. 11); Rho *et al.*, 1969 (p. 158, pl. 1, fig. 6); Hoshino, 1971 (p. 23, pl. 1, fig. 1); Rho & Sim, 1972 b (p. 127); Rho & Lee, 1976 (p. 100).

Material examined: Haeundae, VIII/1982.

Distribution: Korea, Japan.

Genus *Lissodendoryx* Topsent, 1892

11. *Lissodendoryx isodictyalis* (Carter, 1882)

Lissodendoryx isodictyalis: Hartman, 1958 (pp. 41-44, pl. 4, fig. 12, text-fig. 11); Little, 1963 (pp. 48-49); Kim *et al.*, 1968 (pp. 40-41, pl. 2, fig. 11, text-fig. 12); Hoshino, 1971 (p. 23); Rho & Sim, 1972 a (p. 184), 1972 b (p. 127); Rho & Lee, 1976 (pp. 101-102); Hoshino, 1981 (pp. 145-147, text-fig. 61).

Material examined: Sŏngsanp'o, VIII/1982; Namhaedo, VII/1983.

Distribution: Korea, North America, Japan, Indian ocean.

Family Tedaniidae Ridley & Dendy, 1887

Genus *Iotrochota* Ridley, 1884

12. *Iotrochota baculifera* Ridley, 1884

Iotrochota baculifera Ridley, 1884 (pp. 435-436, pl. 39, fig. M, pl. 42, fig. f); Burton, 1959 (p. 239); Tanita, 1969 (pp. 73-74, pl. 2, fig. 6, Text-fig. 1); Rho & Sim, 1976 (p. 74, pl. 7, figs. 3-4); Hoshino, 1976 (p. 6, pl. 2,

figs. 10-12), 1981 (pp. 144-145, pl. 10, figs. 5-6).

Material examined: Söngsanp'o, VI/1984.

Distribution: Korea, Japan, Indian ocean.

Family Haliclonidae De Laubenfels, 1932

Genus *Haliclona* Grant, 1835

13. *Haliclona permollis* (Bowerbank, 1866)

Isodictya permollis Bowerbank, 1866 (p. 278).

Haliclona permollis: De Laubenfels, 1954 (pp. 67-69, text-fig. 38); Tanita, 1958 (p. 130, pl. 1, figs. 3-4, text-fig. 2), 1967 (p. 113), 1968 (p. 41), 1969 (p. 71); Rho *et al.*, 1969 (p. 154); Hoshino, 1974 (p. 8, pl. 1, fig. 2).

Material examined: Sangju-ri, VII/1983; Mijo-ri, VII/1983; Tölsando, VIII/1987.

Distribution: Cosmopolitan.

Family Callyspongidae De Laubenfels, 1936

Genus *Callyspongia* Duchassaing & Micheloffi, 1864

14. *Callyspongia elegans* (Thiele, 1899)

Spinoseella elegans Thiele, 1899 (pp. 23-24, pl. 3, fig. 2, pl. 5, fig. 19).

Callyspongia elegans: Tanita, 1965 (pp. 46-47, pl. 1, fig. 2); Kim *et al.*, 1968 (p. 38, pl. 1, fig. 2, text-fig. 3); Rho *et al.*, 1969 (p. 154); Tanita, 1970 (p. 101); Sim, 1982 (pp. 196-197).

Material examined: Pijin do, VII/1984; Sögwip'o, VII/1987.

Distribution: Korea, Japan, Celebes.

15. *Callyspongia elongata* (Ridley & Dendy, 1886)

Callyspongia elongata: Tanita, 1964 (p. 17, pl. 1, fig. 3), 1967 (p. 114), 1968 (p. 42), 1969 (p. 73); Rho *et al.*, 1969 (p. 155, pl. 1, fig. 1, text-fig. 1); Rho & Sim, 1972a (p. 183), 1972b (p. 125); Rho & Lee, 1976 (pp. 104-105); Sim, 1982 (p. 196).

Material examined: Sögwip'o, VII/1985.

Family Renieridae Schmidt, 1870

Genus *Gellius* Gray, 1867

16. **Gellius edaphus* De Laubenfels, 1930

(pl. 2, figs. 3-4)

Gellius edaphus De Laubenfels, 1932 (p. 111, fig. 66).

Material examined: 33°00'00"N, 127°30'00"E, 145 depth, IV/1987.

Remarks: Sponge is small fragment, 4.5 × 2cm Texture is a little hard, colour in spirit is dirty grey.

Spicules:

Oxea 210-326 × 1 – 18µm.

Sigma 36-52µm.

Distribution: California, Korea.

Family Spongiidae Gray, 1867

Genus *Spongia* Linné, 1759

17. *Spongia officinalis* Linné, 1759

Spongia officinalis: Burton, 1934 (pp. 576-577); De Laubenfels, 1948(p. 4, pl. 1, figs. 1-2); Sim, 1985 (p. 4, pl. 1, figs. 1-2).

Material examined: Söngsanp'o, VII/1984.

Distribution: Korea, Mediterranean Sea, West Indian ocean, South America, Asia.

Famil Tethyidae Gray, 1867

Genus *Tethya* Lamark, 1815

18. *Tethya amamensis* Thiele, 1898

Tethya amamensis Thiele, 1898 (p. 30, pl. 7, figs. 19a-f); Tanita, 1969 (p. 77, pl. 2, fig. 8); Hoshino, 1971 (p. 21); Rho & Sim, 1972a (pp. 185-186, pl. 5, figs. 11-14); 1972b (p. 129).

Material examined: Söngsanp'o, VII/1982.

Distribution: Korea, Japan.

Family Suberitidae Ridley & Dendy, 1886

Genus *Suberites* Nardo, 1833

19. *Suberites excellens* (Thiele, 1898)

Rhizaxinella excellens Thiele, 1898 (p. 34, pl. 3, fig 2, pl. 8 figs. 2a-e).

Suberites excellens: Rho & Lee, 1976 (p. 96, pl. 1, figs. 1-2).

Material examined: Anhuŋg, X/1984.

Distribution: Korea, Japan.

Family Tetillidae Sollas, 1886

Genus *Tetilla* Schmidt, 1868

20. *Tetilla ovata* Thiele, 1898

Craniella ovata Thiele, 1898 (p. 27, pl. 5, fig. 16, pl. 7, figs. 15a-e).

Tethya ovata: Lendenfeld, 1903 (p. 24).

Tetilla ovata: Tanita, 1965 (p. 51, pl. 2, fig. 10), 1969 (p. 76); Rho & Sim, 1972b (p. 130, pl. 4, figs. 1-4).

Material examined: Ch'u ja do, II/1986.

Distribution: Korea, Japan.

Family Raspailiidae Hentschel, 1912

Genus *Raspailia* Nardo, 1833

21. *Raspailia hirsuta* Thiele, 1898

Raspailia hirsuta Thiele, 1898 (p. 59, pl. 3, fig. 9, pl. 8, figs. 46a-d); Tanita, 1970 (p. 102, pl. 2, fig. 8); Rho & Sim, 1972a (p. 185, pl. 3, figs. 7-8); Hoshino, 1977 (p. 6); Sim, 1982 (p. 200, pl. 2, fig. 2).

Material examined: Chejudo, VII/1984.

Distribution: Korea, Japan.

Family Kaliapsidae De Laubenfels, 1936

Genus *Discodermia* Bocage, 1869**22. *Discodermia calyx* Döderlein, 1883**

Discodermia calyx Döderlein, 1883 (p. 77, pl. 5, figs. 4-5); Tanita, 1970 (p. 102, pl. 2, fig. 12); Rho & Sim, 1972b (p. 129, pl. 3, figs. 6-9); Hoshino, 1975 (p. 34, pl. 4, figs. 5-6), 1977 (p. 6).

Material examined: Off Sangju, I/1983, Sögwip'o VII/1984.

Distribution: Korea, Japan.

Family Stelletidae Carter, 1875

Genus *Stelletta* Schmidt, 1862

23. **Stelletta validissima* Thiele, 1898 (pl. 5, figs. 1-2)

Stelletta validissima Thiele, 1898 (p. 13, pl. 1, fig. 5, pl. 7, figs. 1a-i).

Material examined: Chujado, II/1986.

Remarks: Sponge is round shape, 4 cm in diameter, with long root tuft. Texture is hard and tough. Colour in spirit is dirty grey.

Spicules:

Oxea	3900-4920 × 21 – 42 μm.
Dichotriaene	rabdomes 3360 – 5760 × 57 – 75 μm. clad 165 – 204 μm.
Protriaene	rabdomes 3900 – 4800 × 2 – 48 μm. clad 100 – 115 μm.
Large anatriaene	rabdomes 3870 – 4500 × 21 – 51 μm. clad 60-90 μm.
Small anatriaene	rabdomes 3500 – 4500 μm. clad 35 – 54 μm.
Chiaster	6 – 12 μm.
Oxyaster	9 – 33 μm.

Material examined: Ch'ujado, II/1986.

Distribution: Japan, Korea.

Family Aphrocallistidae Gray, 1858

Genus *Aphrocallistes* Gray, 1858

24. **Aphrocallistes jejuensis*, nov. sp.

(pl. 6, figs. 1-3, pl. 7, figs. 1-4, pl. 8, fig. 1, pl. 9, fig. 1, pl. 10, fig. 1)

Material examined: Holotype (Por. 5), Sögwip'o (Pömsö'm), 60m, 11/1971, B.J. Rho. Paratype (Por. 5-1, Por. 5-2), same data as holotype. The type specimens are deposited in the Department of Biology, Ewha Womans University.

Description: This sponge from is three side branched round tubes, 2.5cm to 3.5cm in width. They don't grow straight upward. The tubes being generally 0.6 - 0.8 mm in diameter are not over 1mm in thickness of the wall. The dictyonal framework forms very irregular meshes. The parietal skeleton is honeycomblake, with hexagonal. The outer surface is covered with the dermal membrane. The gastral skeleton is without scopulae. In the dermal skeleton occur hexacts with a variously developed distal

ray, which is bruch type. The proximal ray equals or usually exceeds the four transvers ray in length. Except the dermal hexacts, three types of dermal scopulae occur: one type presents terminally rounded stalk, the other type two or four strong, barbed, unknobbed prongs, the third type two or four straight pointed prongs. The gastral skeleton is without scopulae; it consists of long rod-like diacts, which are rough throughout and are provided with a central node of intersection. The uncinates of the parenchyma vary greatly in length and form. The numerous irregularly scattered hexasters are discohexasters in which the terminal rays are also curved, and they are provided with rounded terminal knobs and have also elongate hemidiscohexaster.

Spicules:

Uncinates	579 - 933 × 3 - 5μm.
Hexact pinulus	106 - 140μm axes ray in length
Leptoscopule	237 - 261μm in total length 60 - 66μm terminal branches
Pachyscopule	387 - 420μm in total length 90μm terminal branches
Scopule with pointed prongs	182 - 196μm total length 28 - 42μm terminal branches
Discohexaster	27 - 55μm in diameter
Hemidiscohexaster	40 - 67μm in diameter

Remarks: This new species is similar to *Aphrocallistes ramosus* (Schulze, 1887) in body form but differs in branches and tube closed with lid which is made of sieve plate. Ijima (1926) and Burton (1959) synonymized *A. ramosus*, *A. bocagei* and *A. beatrix orientalis* with *A. beatrix*; all of these species have oxyhexasters and discohexasters. The new species has discohexasters, hemidiscohexasters and scopulae with pointed prongs, but no oxyhexasters.

ABSTRACT

The sponges identified in the present study turned out to be twenty three species of Demospongia and one Hexactinellida. Of which, six species are new to the Korean fauna and a new species, *Aphrocallistes jejuensis* belonging to Aphrocallistidae, Hexasterophora was found from pömsö, cheju-do.

REFERENCES

- Bowerbank, J. S., 1866. A monograph of the British spongiidae II. Ray. Soc. Publ. pp. 1-388.
 Burton, M., 1934. Sponges. Great Barrier Reef Expedition (1928-29) Report. British Mus. Nat. Hist., 4, 14: 513-614.
 Burton, M., 1959. Sponges. The John Murray Expedition 1933-34 Sci. Rep., 10, 5: 150-281.
 De Laubenfels, M. W., 1932. The Marine and Fresh-Water Sponges of California. U.S. Nat. Mus., 81 (2927): 1-140.
 De Laubenfels, M. W., 1935. Some Sponges of Lower California (Mexico). American Museum Novitates. No. 779: 1-14.
 De Laubenfels, M. W., 1948. The Order Keratosa of the Phylum Porifera—A Monographic Study. Allan Hancock Foundation Occ., 3: 1-195.


- De Laubenfels, M. W., 1954. The Sponges of the West-Central Pacific. *Studies in Zool.*, **7**: 1-319.
- Dendy, A., 1921. Report on the Sigmatotetraxonida Collected by H.M.S. "Sealark" in the Indian ocean. *Trans. Linn. Soc. Ser. 2 Zool.*, **18**: 1-164.
- Döderlein, L., 1883. Studien an Japanischen Lithistiden. *Zeit. f. Wiss. Zool.*, **40**: 62-104.
- Hartman, W. D., 1958. Natural History of the Marine Sponges of southern New Zealand. *Peabody Mus. Nat. Hist. yale Univ. Bull.*, **12**: 1-155.
- Hoshino, T., 1970. 15 Sponges (Demospongiae) obtained from Mukaishima and Its Adjacent Waters. *Bull. Biol. Soc. Hiroshima Univ.*, 36-21-26.
- Hoshino, T., 1971. Sponge Fauna of Seto Inland Sea (Demospongiae, Calcarea). *Bull. Biol. Soc. Hiroshima Univ.*, **38**: 21-30.
- Hoshino, T., 1974. Demospongia of Hiryu-Jima (Biro-Jima), an Islet in the Ariake Sea. *Calanus.*, **4**: 8-15.
- Hoshino, T., 1975. The Sponges of the Anan Coast. *Zool. Mag.*, **84**, 1:30-38.
- Hoshino, T., 1976. Demosponges Obtained from the Vicinity of the Aitsu Marine Biological Station. *Calanus*, **5**: 3-11.
- Hoshino, T., 1977. Demosponges from the Kii Channel and its Environs Western Japan. *Proc. Jap. Soc. Syst. Zool.*, **13**: 5-15.
- Hoshino, T., 1981. Shallow-Water Demosponges of Western japan, I. *Jour. Sci. Hiroshima Univ. Ser. B. Div. 1 (Zool.)*, **29**, 1: 47-183.
- Ijima, I., 1926. The Heactinellida of the Siboga Expedition. *Siboga Exped. Rep. Leiden.*, Vol. CVI, pp. 1-383.
- Kim, H. S., B. J. Rho and C. J. Sim, 1968. Marine Sponges in South Korea (I). *Korean Jour. Zool.*, **11**, 2: 37-47.
- Lendenfeld, R. Von., 1903. Tetraxonida. *Das Tierreich* **19**: 1-168.
- Little, F. J. Jr., 1963. The Sponge Fauna of the St. George's Sound, Apalachee Bay, and Panama City Regions of the Florida Gulf Coast. *Tulane Studies in Zool.*, **11**, 2: 31-71.
- Rho, B. J., H. S. Kim and C. J. Sim, 1969. Marine Sponges in South Korea (2). *Jour. Kor. Res. Inst. Bet. Liv.*, **3**: 153-160.
- Rho, B. J. and C. J. Sim, 1972b. Faunal Studies on the Sponges in Korea. R-72-82, *Min. Sci. Tech.*, 121-138.
- Rho, B. J. and C. J. Sim, 1976. On the Classification and the Distribution of the Marine Benthic Animals in Korea *Jour. Res. Inst. Bet. Liv.*, **16**: 67-87.
- Rho, B. J. and K. H. Lee, 1976. On the classification and the Distribution of the Marine Benthic Animals in Korea (4). Sponges. *Jour. Res. Inst. Bet. Live.*, **16**: 67-87.
- Ridley, S. O., 1884. Spongiida, Report on the Zoological Collections Made in the Indo-Pacific Ocean during the Voyage of the H.N.S. "Alert" 1881-82. *Brit. Mus. Hist.* : 366-482.
- Schulze, F. E., 1887. Report on the Hexactinellida collected by H.M.S. Challenger during the years 1873-1876. *Rep. Scient. Result Challenger. Zoology.*, Vol. XII, part LIII, PP. 1-514, pl. I-CIV.
- Sim, C. J., 1982. A Systematic Study on the Marine Sponges from Jeju Island. *Soong Jun. Univ. Eassays and Papers*, **12**: 187-210.
- Sim, C. J., 1985. A Systematic Study on the Marine Sponges from the South Sea and the Yellow sea of Korea. *Korean J. Syst. Zool.*, **1**, 1-2: 21-30.
- Tanita, S., 1958. Sponges Collected from Oxyter-Rafts in Matsushima Bay and Its Adjacent Waters. *Bull. Tohoku Reg. Fish. Res. lab.*, **11**: 127-143.
- Tanita, S., 1961. Report on the sponges Collected from the Kurushima Strait, Seto Inland Sea. *Mem. Ehime Univ. Sect. II. Ser. B.*, **4**, 2: 151-170.
- Tanita, S., 1963. Report on the Non-Calcareous Sponges in the Museum of the Biological Institute of Tohoku University, Part II. *Sci. Rep. Tohoku Univ. Ser. 4. Biol.*, **29**: 121-129.
- Tanita, S., 1964. Sponges Obtained from Tsukumo Bay and Its Vicinity. *Ann. Rep. Noto Mar. lab.*, **4**: 15-22.

- Tanita, S., 1965. Report on the Sponges obtained from the Adjacent Waters of the Sado Island, Japan Sea. Bull. Jap. Sea. Reg. Fish. Res. Lab., **14**: 43-66.
- Tanita, S., 1969. Further Studies on the Sponges Obtained from the Sado Island and Its Adjacent Waters. Bull. Jap. Sea. Reg. Fish. Res. Lab., **21**: 67-88.
- Tanita, S., 1970. The Sponges in the Tokushima Musium. Bull. Tohoku Reg. Fish. Res. Lab., **30**: 99-105.
- Thiele, J., 1898. Studien über Pazifische Spongie. Zoologica, **24**: 1-72.
- Thiele, J., 1899. Studien über Pazifische Spongien. *ibid.*, **24**: 5-33.

RECEIVED: 13 FEBRUARY 1988

ACCEPTED: 3 MAY 1988


PLATE 1



Figs. 1-2. *Ophlitaspongia minor* Burton.

1. Entire animal; 2. Megasclere: A, Large Style; B, Small Style; C, Subtylostyle
Microsclere: D, Large toxa; E, Small toxa; F, Isochela.

PLATE 2





Figs. 1-2. *Clathria madrepora* Dendy.

1. Entire animal; 2. Megasclere: A, Large tylostyle; B, Small tylostyle; C, Acanthostyle.

Figs. 3-4. *Gellius edaphus* De Laubenfels.

3. Entire animal; 4. Megasclere: A, Oxea, Microsclere: B, Sigma.


PLATE 3



Figs. 1-2. *Myxilla productus* Hoshino.

1. Entire animal; 2. Megasclere: A, Tornote; B, Large acanthostyle; C, Small acanthostyle.
 Microsclere: D, Large isochela; E, Sigma; F, Small isochela.

PLATE 4




Figs. 1-2. *Oxymycale paradoxa* De Laubenfels.

1. Entire animal; 2. Megasclere: A, Oxea; B, Style.

Microclere: C, Anisochela; D, Sigma; E, Raphid.


PLATE 5



Figs. 1-2. *Stelletta validissima* Thiele.

1. Entire animal; 2. Megasclere: A, Dichotriaene; B, Large anatriaene; C, Small anatriaene; D, Protriaene; E, Oxea. Microsclere: F, Oxyaster; G, Chiaster.


PLATE 6



Figs. 1-3. *Aphrocallistes jejuensis*, nov. sp.

1. Entire animal; 2. Surface; 3. Dictyonal skeleton.

PLATE 7



Figs. 1-4. *Aphrocallistes jejuensis*, nov. sp.

1. Scopule; 2. Gastral part; 3-4. Surface with scopules.




Fig. 1. *Aphrocallistes jejuensis*, nov. sp.

1. Megascclere: A-B, Hexact; C, Uncinate; D, Hexact.

PLATE 9





Fig. 1. *Aphrocallistes jejuensis*, nov. sp.

1. Microsclere: A, Pachyscopule; B, Leptoscopule; C, Scopule with pointed prongs.

PLATE 10



Figs. 1-2. *Aphrocallistes jejuensis*, nov. sp.

1. Microsclere: A-B, Heact pinulus; 2. Microsclere: A, Dischohexaster; B, Hemidischohexaster.